

Application Advice

Hops

Standard application rates for organic farming

Organic agriculture As of May 2024



Application rate | per ha

Spreading

Soil fertilising

AKRA KOMBI	250–300 kg
AKRA DGC	Application depending on fractional soil analyses

Application rate for planting out – seedlings (saplings)

Water	100 l
AKRA Saat-Power	1 l

- ⚙ Leave seedlings in the solution for 2–3 hours
- ⚙ Leave potted plants in the solution for 2 minutes

Spraying

At vegetation start

Water	350 l
AKRA Stroh R+P+K	2 l
AKRA MSB	2 l
AKRA Sulpur+	2 l
AKRA WD	2 l
Wetting agent	

With shoot length of 30 cm

Water	400 l
AKRA Plus 9	0.5 l
AKRA Blatt	2 l
AKRA MSB	0.5 %
AKRA Sulpur+	0.2 %
AKRA WD	0.1 %

With shoot length of 100cm

Water	600–1000 l
AKRA Plus 9	0.2 %
AKRA Blatt	0.8 %
AKRA MSB	0.5 %
AKRA Azotobacter	0.1 %
AKRA Sulpur+	0.3 %
AKRA WD	0.2 %

With a shoot length from 200 cm (when needed)

Water	1000–1200 l
AKRA Plus 9	0.2 %
AKRA Blatt	0.8 %
AKRA MSB	0.5 %
AKRA Azotobacter	0.2 %
AKRA Sulpur+	0.3 %
AKRA WD	0.2 %

INFO

- ⚙️ Seek specific information when dealing with severe infestations of spider mites, rust mites, red spider mites or aphids.
- ⚙️ During bud spraying, apply the solution over the entire area.
- ⚙️ The pH value of the spray solution should be adjusted to 5.5 with citric acid.
- ⚙️ The addition of **AKRA** foliar fertiliser can and may reduce the pH value of the spray solution.
- ⚙️ Mix in any other products individually once the pH value has been adjusted.
- ⚙️ Make sure that you add the additional products in the right order.
- ⚙️ The dosage of **AKRA** foliar fertiliser must be adjusted to the quantity of water used.

SUCCESS

WITH THE **AKRA FERTILISING SYSTEM**

» Guaranteed

Page: 3/3



retrieve application
advice online

