


Application Advice

Poppy

Conventional agriculture As of May 2024



Stages of development



1. Seed treatment | for 100 kg of seeds

AKRA Saat-Power	0.4–0.5 l
-----------------	-----------

2. Spraying (cotyledon stage / 1st foliage leaf)

WARNING Watch out for flea beetles!

Water	150 l
AKRA Plus 9	0.25l
AKRA Blatt	0.5 l
AKRA MSB	0.5 l
AKRA Sulpur+	0.75 l
AKRA WD	0.25 l

3. Spraying (start of shoot formation)

Water	200 l
AKRA Plus 9	0.5 l
AKRA Blatt	2 l
Fungicide	When needed (50 % of the recommended application rate)
AKRA MSB	1 l
AKRA Azotobacter	0.25 l
AKRA Sulpur+	1 l
AKRA WD	0.5 l
Urea	5 kg
Wetting agent	

4. Spraying (before flowering)

Water	200 l
AKRA Plus 9	0.5 l
AKRA Blatt	2 l
Fungicide	When needed (50 % of the recommended application rate)
AKRA MSB	1 l
AKRA Azotobacter	0.5 l
AKRA Sulpur+	2 l
AKRA WD	1 l
Urea	10 kg
Insecticide	Depending on infestation pressure
Wetting agent	

5. Spraying (green capsule stage)

Water	300 l
AKRA Plus 9	1 l
AKRA Blatt	3 l
AKRA MSB	2 l
AKRA Azotobacter	1 l
AKRA Sulpur+	3 l
AKRA WD	2 l
Wetting agent	1 ml per 100 l of solution

INFO

- ⚙ The pH value of the water should be adjusted to 5.5 with citric acid.
- ⚙ The addition of **AKRA** foliar fertiliser can and may reduce the pH value of the water.
- ⚙ Mix in any other additives once the pH value has been adjusted.
- ⚙ Make sure that you add the additional products in the right order.
- ⚙ Applications 4 and 5 should be sprayed around midday.
- ⚙ The dosage of **AKRA** foliar fertiliser must be adjusted to the quantity of water used.

SUCCESS

WITH THE **AKRA FERTILISING SYSTEM**

» Guaranteed

