Application Advice






Standard application rates for organic farming

Organic agriculture As of May 2024



Stages of development





1. Seed treatment | for 100 kg of seeds

AVDA Cook Downey	0.4.0.5.1
AKRA Saat-Power	0.4-0.51

2. Spraying (cotyledon stage / 1st foliage leaf)

WARNING Watch out for flea beetles!

Water	150 l
AKRA Plus 9	0.25l
AKRA Blatt	0.5
AKRA MSB	0.5
AKRA Sulpur+	11
AKRA WD	0.25

Application Advice **Poppy**



3. Spraying (start of shoot formation)

Water	200
AKRA Plus 9	0.5
AKRA Blatt	21
AKRA MSB	11
AKRA Azotobacter	0.25 l
AKRA Azo+	0.25 l
AKRA Sulpur+	2
AKRA WD	11
AKRA org. liquid fertiliser	20
Wetting agent	

4. Spraying (before flowering)

Water	200 l
AKRA Plus 9	0.5
AKRA Blatt	21
AKRA MSB	11
AKRA Azotobacter	0.5
AKRA Sulpur+	31
AKRA WD	1
AKRA org. liquid fertiliser	20
Wetting agent	

Application Advice Poppy



5. Spraying (green capsule stage)

Water	300
AKRA Plus 9	11
AKRA Blatt	31
AKRA MSB	31
AKRA Azotobacter	11
AKRA Sulpur+	5 l
AKRA WD	2
Wetting agent	

INFO

- The pH value of the water should be adjusted to 5.5 with citric acid.
- The addition of **AKRA** foliar fertiliser can and may reduce the pH value of the spray solution.
- Mix in any other additives once the pH value has been adjusted.
- Make sure that you add the additional products in the right order.
- Applications 4 and 5 should be sprayed around midday.
- The dosage of AKRA foliar fertiliser must be adjusted to the quantity of water used.



