

# Application Advice Strawberries

Field-grown

Conventional and organic agriculture As of May 2024



## Planting out and Spreading

### Application rate for planting out

Water	100 l
<b>AKRA</b> Saat-Power	1 l

- ⚙ Leave seedlings in the solution for 1–2 hours (for bare root plants)
- ⚙ Leave potted plants in the solution for 2 minutes

### Spreading

<b>AKRA</b> Kombi	500 kg annually
<b>AKRA</b> DGC	As needed depending on fractional soil analyses
Nitrogen (urea)	As needed

## Liquid fertiliser application (also for organic farms)

### Bud spraying at vegetation start

Water	300 l
<b>AKRA</b> MSB	3 l
<b>AKRA</b> Sulpur	3 l
<b>AKRA</b> WD	3 l
Wetting agent	

## Start of flowering

Water	200 l
<b>AKRA Plus 9</b>	0.5 l
<b>AKRA Blatt</b>	2 l
<b>AKRA Azotobacter</b>	0.5 l
Wetting agent	

## Before fruit turns red

Water	250–300 l
<b>AKRA Plus 9</b>	1.5 l

## INFO

- ⚙ The pH value of the spray solution should be adjusted to 5.5 with citric acid.
- ⚙ The addition of **AKRA** foliar fertiliser can and may reduce the pH value of the spray solution.
- ⚙ Mix in any other products individually once the pH value has been adjusted.
- ⚙ Make sure that you add the additional products in the right order.
- ⚙ The dosage of **AKRA** foliar fertiliser must be adjusted to the quantity of water used.
- ⚙ When using fungicide, only use 50 % of the recommended application rate.
- ⚙ Use insecticide as needed.
- ⚙ **Follow AKRA's advice on soil preparation for new planting.**
- ⚙ **Treat the harvest residue in autumn ONCE THE LEAVES HAVE BEEN CUT!**

**SUCCESS**

WITH THE **AKRA FERTILISING SYSTEM**

» Guaranteed

